

TABEL. *Overzicht van de prevalentie van neurologische en psychiatrische klachten in een cohort van 236379 ex-Covid-19-patiënten in de VS.^[3]*

	All patients	Patients without hospitalisation	Patients with hospitalisation	Patients with ITU admission	Patients with encephalopathy
Intracranial haemorrhage (any)	0.56% (0.50-0.63)	0.31% (0.25-0.39)	1.31% (1.14-1.52)	2.66% (2.24-3.16)	3.61% (2.97-4.39)
Intracranial haemorrhage (first)	0.28% (0.23-0.33)	0.14% (0.10-0.20)	0.63% (0.50-0.80)	1.05% (0.79-1.40)	1.19% (0.82-1.70)
Ischaemic stroke (any)	2.10% (1.97-2.23)	1.33% (1.22-1.46)	4.38% (4.05-4.74)	6.92% (6.17-7.76)	9.35% (8.23-10.62)
Ischaemic stroke (first)	0.76% (0.68-0.85)	0.43% (0.36-0.52)	1.60% (1.37-1.86)	2.82% (2.29-3.47)	3.28% (2.51-4.27)
Parkinsonism	0.11% (0.08-0.14)	0.07% (0.05-0.12)	0.20% (0.15-0.28)	0.26% (0.15-0.45)	0.46% (0.28-0.78)
Guillain-Barré syndrome	0.08% (0.06-0.11)	0.05% (0.03-0.07)	0.22% (0.15-0.32)	0.33% (0.21-0.54)	0.48% (0.20-1.14)
Nerve, nerve root, or plexus disorders	2.85% (2.69-3.03)	2.69% (2.51-2.89)	3.35% (3.02-3.72)	4.24% (3.58-5.03)	4.69% (3.81-5.77)
Myoneural junction or muscle disease	0.45% (0.40-0.52)	0.16% (0.12-0.20)	1.24% (1.05-1.46)	3.35% (2.76-4.05)	3.27% (2.54-4.21)
Encephalitis	0.10% (0.08-0.13)	0.05% (0.03-0.08)	0.24% (0.17-0.33)	0.35% (0.19-0.64)	0.64% (0.39-1.07)
Dementia	0.67% (0.59-0.75)	0.35% (0.29-0.43)	1.46% (1.26-1.71)	1.74% (1.31-2.30)	4.72% (3.80-5.85)
Mood, anxiety, or psychotic disorder (any)	23.98% (23.58-24.38)	23.59% (23.12-24.07)	24.50% (23.76-25.26)	27.78% (26.33-29.29)	36.25% (34.16-38.43)
Mood, anxiety, or psychotic disorder (first)	8.63% (8.28-8.98)	8.15% (7.75-8.57)	8.85% (8.22-9.52)	12.68% (11.28-14.24)	12.96% (11.13-15.07)
Mood disorder (any)	13.66% (13.35-13.99)	13.10% (12.73-13.47)	14.69% (14.09-15.32)	15.43% (14.27-16.68)	22.52% (20.71-24.47)
Mood disorder (first)	4.22% (3.99-4.47)	3.86% (3.60-4.14)	4.49% (4.05-4.99)	5.82% (4.86-6.97)	8.07% (6.56-9.90)
Anxiety disorder (any)	17.39% (17.04-17.74)	17.51% (17.09-17.93)	16.40% (15.76-17.06)	19.15% (17.90-20.48)	22.43% (20.65-24.34)
Anxiety disorder (first)	7.11% (6.82-7.41)	6.81% (6.47-7.16)	6.91% (6.38-7.47)	9.79% (8.65-11.06)	9.24% (7.70-11.07)
Psychotic disorder (any)	1.40% (1.30-1.51)	0.93% (0.83-1.04)	2.89% (2.62-3.18)	2.77% (2.31-3.33)	7.00% (6.01-8.14)
Psychotic disorder (first)	0.42% (0.36-0.49)	0.25% (0.19-0.33)	0.89% (0.72-1.09)	0.70% (0.46-1.06)	2.12% (1.53-2.94)
Substance use disorder (any)	6.58% (6.36-6.80)	5.87% (5.63-6.13)	8.56% (8.10-9.04)	10.14% (9.25-11.10)	11.85% (10.55-13.31)
Substance use disorder (first)	1.92% (1.77-2.07)	1.74% (1.58-1.91)	2.09% (1.82-2.40)	3.15% (2.60-3.82)	2.58% (1.91-3.47)
Insomnia (any)	5.42% (5.20-5.64)	5.16% (4.91-5.42)	5.95% (5.53-6.39)	7.50% (6.66-8.44)	9.82% (8.57-11.24)
Insomnia (first)	2.53% (2.37-2.71)	2.23% (2.05-2.43)	3.14% (2.81-3.51)	4.24% (3.55-5.07)	5.05% (4.10-6.20)
Any outcome	33.62% (33.17-34.07)	31.74% (31.22-32.27)	38.73% (37.87-39.60)	46.42% (44.78-48.09)	62.34% (60.14-64.55)
Any first outcome	12.84% (12.36-13.33)	11.51% (10.98-12.07)	15.29% (14.32-16.33)	25.79% (23.50-28.25)	31.13% (27.29-35.36)

Data are percentage at 6 months (95% CI). Additional outcomes are presented in the appendix (pp 27-28). ITU=intensive therapy unit.